



CONFIDENTIAL SEXUALLY TRANSMITTED DISEASE (STD) CASE REPORT

Report STDs within three working days. (WAC 246-101-101/301)

PATIENT INFORMATION
Last Name, First Name, Middle Initial, Date of Birth
Address, City, State, Zip Code
Email Address, Telephone
Reason for Exam (check one)
Date of Diagnosis, Sex, If Female, Pregnant?
Ethnicity, Race (check all that apply), Gender of Sex Partners, HIV Tested at This Visit?
DIAGNOSIS—DISEASE
GONORRHEA (Lab Confirmed), CHLAMYDIA TRACHOMATIS (Lab Confirmed)
SYPHILIS, HERPES SIMPLEX
PARTNER MANAGEMENT PLAN—Select method of ensuring partner treatment
REPORTING CLINIC INFORMATION

Thank you for reporting a STD. All information will be managed with the strictest confidentiality.

PRIVILEGED AND CONFIDENTIAL COMMUNICATION: The information contained in this message is privileged, confidential or otherwise exempt from disclosure and is intended solely for the use of the individual(s) named above. If you are not the intended recipient, you are hereby advised that any dissemination, distribution or copying of this communication is prohibited. If you have received this facsimile in error, please immediately notify the sender by telephone and destroy the original facsimile. (8/2016) Page 1

PARTNER MANAGEMENT PLAN INSTRUCTIONS

| |
|---|
| PARTNER TREATMENT |
| <p>Chlamydia & Gonorrhea Providers are to manage partner treatment by either treating partners or prescribing medication.</p> <p>All partners should be treated as if they are infected.</p> <ul style="list-style-type: none"> • Examine and treat all the patient’s sex partners from the previous 60 days. If this is not possible, offer medication for all sex partners whom patients are able to contact. <p>Cowlitz County Health Department investigates ALL Syphilis cases, and certain Chlamydia & Gonorrhea cases (patients <16 years old, or men having sex with men).</p> |
| OTHER STDS: PARTNER TREATMENT |
| <p>Genital herpes, chancroid, granuloma inguinale, or lymphogranuloma venereum diagnosis Health Department does not follow-up with cases. Advise patient to notify sex partners. Partners should contact their provider for testing and treatment.</p> |

RECOMMENDED REGIMENS FOR ANTIMICROBIALS LISTED ON CASE REPORTS*

| |
|--|
| GONORRHEA—UNCOMPLICATED |
| <p>Ceftriaxone 250 mg IM as a single dose PLUS Azithromycin 1g PO as a single dose</p> <p>Alternatives: Cefixime 400 mg PO as a single dose..... PLUS Azithromycin 1g PO as a single dose</p> <p>For beta-lactam allergic patients: Azithromycin....2g PO as a single dose....PLUS Gentamicin 240mg IM as a single dose OR Gemifloxacin 320mg PO as a single dose</p> <p><i>Fluoroquinolones (Levofloxacin or Ciprofloxacin, etc.) are no longer recommended for the treatment of gonorrhea due to increased prevalence of quinolone-resistant Neisseria gonorrhoeae (QRNG).</i></p> |
| CHLAMYDIA—UNCOMPLICATED |
| <p>Azithromycin..... 1g PO as a single dose</p> <p>OR</p> <p>Doxycycline 100 mg PO BID for 7 days (Preferred for rectal chlamydia infection)</p> <p>Alternatives: Erythromycin (base) 500 mg PO QID for 7 days OR Ethylsuccinate..... 800 mg PO QID for 7 days OR Ofloxacin 300 mg PO BID for 7 days OR Levofloxacin 500 mg PO for 7 days</p> |
| SYPHILIS—PRIMARY, SECONDARY OR EARLY LATENT (<1 YEAR) |
| <p>Benzathine penicillin G 2.4 million units IM in a single dose</p> |
| SYPHILIS—LATE LATENT, LATENT OF UNKNOWN DURATION, TERTIARY (NOT NEUROSYPHILIS) |
| <p>Benzathine penicillin G 2.4 million units IM for 3 doses at 1 week intervals</p> |

* Refer to "STD Diagnostic and Treatment Guidelines" or the Centers for Disease Control and Prevention's (CDC's) website (www.cdc.gov/std/treatment) for further information on treating pregnant patients, infections of the pharynx, treatment of infants and other details.