

# Poison Oak

## Integrated Pest Management Control Measures

### Mechanical

CAUTION!! Use protective clothing and gloves when handling this plant due to skin irritants. Mowing young shoots is effective, but will only provide partial control. Cutting or digging can be effective. Remove entire plant and root system in early spring or late summer for best results.

### Cultural

Goats and sheep will easily graze on poison oak without side effects, but both animals are not selective and will graze on other vegetation on site.

### Biological

No biological controls are currently available in the United States.

### Chemical

*Always read herbicide label before applying*

*Toxicodendron diversilabum*



FOLIAR SPRAY			
Chemical	Brand Names	Label Use	Calendar/Plant Stage
2,4D	Weedmaster	Cropland, pastures, rangeland	Late summer or fall to actively growing plant
triclopyr	Garlon 3A	Non-cropland	Spring to fall when plant is actively growing
triclopyr ester + 2,4D	Crossbow	Rangeland, permanent grass pastures, non-cropland	Spring when plants are actively growing
dicamba	Banvel Clarity	Cropland, pasture, rangeland, turf and grass	Spring and fall when plants are actively growing
triclopyr + 2,4D	PastureGard	Rangeland, permanent pastures	Spring to actively growing plants and after leaves have fully extended and terminal growth has slowed
glyphosate	RoundUp	Non-cropland	
<b>Non Selective Herbicide</b>			



**Cowlitz County Noxious Weed Control Board**  
 Administration Annex Building, 1st Floor ■ Kelso, WA 98626  
 Phone: 360-577-3117 ■ Fax: 360-425-7760  
 E-mail: [velazqueza@co.cowlitz.wa.us](mailto:velazqueza@co.cowlitz.wa.us)